

REJUVenate™

Ready to Use

Efficacy Claims & Use Directions



One-Step Disinfectant Cleaner FOR USE IN SPAS, SALONS & CLINICS

**KEEP OUT OF REACH
OF CHILDREN**

READ S.D.S. BEFORE
USING THE PRODUCT
S.D.S. available online at
www.virox.com

Manufactured by/for:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON CANADA L6H 6R1
1-800-387-7578 | virox.com

Meets OSHA Bloodborne Pathogen Standard for *HBV, *HCV & *HIV-1 | Bactericidal, *Virucidal, Pseudomonacidal, Fungicidal, and Tuberculocidal

For Use in Spas, Salons & Clinics. Ideal for Pedicure Foot Spas

Rejuvenate™ Ready to Use One-Step Disinfectant Cleaner is designed for cleaning and disinfection of hard non-porous surfaces in beauty salons, tanning facilities, spas, nail salons, tattoo parlors, barbershops, and hairstyling establishments. It cleans by removing body oils, dead tissue, soil and all other buildups of organic matter found on hard non-porous surfaces.

It is designed for use on hard, non-porous environmental surfaces such as foot baths, foot spas, pedicure foot spa equipment and esthetic tools and equipment.

BACTERICIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces

Acinetobacter baumannii

Enterococcus faecalis (VRE)

Escherichia coli O157:H7

Escherichia coli (ESBL)

Klebsiella pneumoniae

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Shigella dysenteriae

Staphylococcus aureus

Staphylococcus aureus (MRSA)

Staphylococcus aureus (Community Associated Methicillin Resistant (CA-MRSA)) (NARSA NRS 384) (Genotype USA 300)

Staphylococcus aureus (Community Associated Methicillin Resistant (CA-MRSA)) (NARSA NRS 123) (Genotype USA 400)

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

* VIRUCIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces

Adenovirus type 8

Avian Influenza A virus

Hepatitis B Virus (HBV)

Hepatitis C Virus (HCV)

Herpes Simplex Virus Type 1 (HSV-1)

Herpes Simplex Virus Type 2 (HSV-2)

HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1)

Human Coronavirus

Influenza A Virus (Strain Hong Kong)

Norovirus (Feline Calicivirus, as the surrogate)

Poliovirus Type 1, Strain Brunhilde

Rhinovirus Type-37, Strain 151-1

Rotavirus (Strain WA)

This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Virucidal activity was determined by the efficacy test methods for virucidal agents intended for inanimate environmental surfaces; ASTM 1053-97 Standard Test Method for Efficacy of Virucidal Agents Intended for Inanimate Environmental Surfaces and EPA protocols for surrogate viral testing.

TUBERCULOCIDAL: 5 MINUTES

In the presence of 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces

[†]*Mycobacterium bovis* (BCG)

Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL: 10 MINUTES

In the presence of 5% serum load and 10 minute contact time on hard, nonporous environmental surfaces

Trichophyton interdigitale (Athlete's Foot and Nail Fungus)

Fungicidal activity was determined by the AOAC Fungicidal Activity of Disinfectants Method.

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 30 SEC

In the presence of 5% serum load and 30 second contact time on hard, non-porous environmental surfaces

Enterococcus faecalis (VRE)

Escherichia coli O157:H7

Klebsiella pneumoniae

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

Staphylococcus aureus (MRSA)

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

SOFT SURFACE[†] SANITIZATION

For use on soft surfaces (blinds, bedding, blankets, chairs, couches, curtains, drapes, linens, mattresses, soft cushions, sheets, sofas, upholstered furniture, wash cloths and window treatments).

[†] composed of cotton or polyester

SOFT SURFACE[†] SANITIZATION AGAINST THE FOLLOWING ORGANISMS IN 1 MINUTE

Enterobacter aerogenes

Staphylococcus aureus

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD/ BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV OR HBV OF SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of Rejuvenate™.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HCV and HBV. Use a 5 minute contact time for Mycobacteria[‡] and a 10 minute contact for fungi.

Disposal of Infectious Material: Cleaning materials used containing blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

Broad Spectrum Non-Food Contact Sanitizer:

Apply to hard non-porous surface; allow to remain wet for 30 seconds. Wipe dry or allow to air dry. No rinsing is required. For heavily soiled areas a pre-cleaning is required.

To Sanitize Soft Surfaces[†]:

Spray a light even coating on soiled fabric until damp. Allow area to remain damp for 1 minute. Gently blot area with a clean, damp, color-safe cloth. Repeat as needed (for stubborn stains or heavy fabrics). Let air dry. To test for color fastness, locate a small inconspicuous area on your fabric and apply the solution as per the use directions. Once dry, evaluate discoloration.

For use as a disinfectant, on hard, non-porous surfaces of non-circulating foot baths (pedicure foot spa equipment):

- 1. Drain:** Drain the water and remove any visible debris.
- 2. Clean:** Thoroughly clean all (hard, non-porous (inanimate) environmental) surfaces with this product, then rinse with water and drain.
- 3. Apply:** Apply this product on all hard non-porous surfaces with a trigger spray.
- 4. Scrub:** Brush or swab thoroughly and allow solution to stand for 1 minute. Use a 5 minute contact time for Mycobacteria[‡] and a 10 minute contact time for fungi.
- 5. Rinse** all surfaces with fresh water and drain.

To Disinfect Non-Critical[‡], Pre-Cleaned Instruments:

- Instruments must be thoroughly pre-cleaned to remove excess organic debris, rinsed, and dried.
- Clean and rinse lumens of hollow instruments before filling with this product.
- Spray all surfaces of instruments with this product until thoroughly wet. Let stand for 1 minute. Use a 5 minute contact time for Mycobacteria[‡] and a 10 minute contact time for fungi.
- Wipe with a clean, damp cloth or paper towel and allow to air dry

[‡] Non-critical medical devices are items that come in contact only with intact skin.

Note: Critical and Semi-critical devices must be followed up by appropriate terminal sterilization/high-level disinfection process.

[†] composed of cotton or polyester

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant Product:

- Pre-clean heavily soiled areas.
- Apply solution by spray, cloth, disposable wipe or mop to hard, non-porous environmental surfaces.
- All surfaces must remain wet for 1 minute. Use a 5 minute contact time for Mycobacteria[‡] and a 10 minute contact time for fungi.
- Wipe surfaces dry or allow to air dry.

NOTICE TO USER: Rejuvenate™ may not be suitable for all items or surfaces. Prolonged exposure to copper, brass, anodized aluminum and some coatings should be avoided. If you are uncertain as to the material composition of your item, confirm with the manufacturer before proceeding or visit the Compatibility section on www.virox.com for a listing of compatible materials.

Si no puede leer en inglés, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto.